

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):





- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 401/06, 233/56, A61K 31/4164, A61P 25/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/23438</b> <b>(43) International Publication Date:</b> 27 April 2000 (27.04.00)
<b>(21) International Application Number:</b> PCT/US99/21461 <b>(22) International Filing Date:</b> 15 October 1999 (15.10.99) <b>(30) Priority Data:</b> 09/173,642 16 October 1998 (16.10.98) US <b>(71) Applicant:</b> SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US). <b>(72) Inventors:</b> WOLIN, Ronald; 16309 Los Rosales Street, San Diego, CA 92127 (US). ROSENBLUM, Stuart, B.; 16 Steven Terrace, West Orange, NJ 07052 (US). AFONSO, Adriano; 10 Woodmere Road, West Caldwell, NJ 07006 (US). <b>(74) Agents:</b> KALYANARAMAN, Palaiyur, S. et al.; Schering-Plough Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> N-(IMIDAZOLYLALKYL)SUBSTITUTED CYCLIC AMINES AS HISTAMINE-H <sub>3</sub> AGONISTS OR ANTAGONISTS		
<b>(57) Abstract</b>  <p>The present invention discloses novel N(imidazolylalkyl)-substituted cyclic amine compounds of Formula (I), wherein R<sub>1</sub> is selected from the group consisting of H,OCO-R<sub>7</sub>, and CO<sub>2</sub>R<sub>7</sub> with R<sub>7</sub> being a substituted or unsubstituted alkyl or aryl, the substituents on said substituted alkyl or aryl being selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic and halogen; R<sub>2</sub> is selected from the group consisting of H, hydroxyl or halogen; G is spacer moiety selected from the group consisting of a C<sub>3</sub>-C<sub>7</sub> straight chain alkyl, a C<sub>2</sub>-C<sub>7</sub> alkene or a C<sub>2</sub>-C<sub>7</sub> alkyne, with said straight chain alkyl, alkene or alkyne being optionally substituted with one or more R<sub>7</sub> groups; and T which represents a cyclic amine is a ring moiety selected from the group consisting of a, b, c or d, wherein n is an integer for 0-3, and R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> can be the same or different with the proviso that any two of said R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> can be bound to the same or different carbon atom of ring T, with said R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> being independently selected from the group consisting of H, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkenyl, substituted or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkynyl, aryl, aralkyl, alkylaryl, cycloalkyl, heterocyclic, C-O-R<sub>8</sub>, C(=O)R<sub>8</sub>, CO<sub>2</sub>R<sub>8</sub>, OC(O)R<sub>8</sub>, NH<sub>2</sub>NHR<sub>8</sub>, N(R<sub>8</sub>)<sub>2</sub>, NR<sub>8</sub>R<sub>9</sub>, SR<sub>8</sub>, SO<sub>2</sub>R<sub>8</sub>, S(O)R<sub>8</sub>, OH,OR<sub>8</sub>, CH<sub>2</sub>OR<sub>8</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>N(R<sub>8</sub>)<sub>2</sub>, CH<sub>2</sub>NHR<sub>8</sub>, CH<sub>2</sub>SR<sub>8</sub>, C(O)-NHR<sub>8</sub>, C(O)NR<sub>8</sub>R<sub>9</sub>, CN,NO<sub>2</sub>, NR<sub>8</sub>-CO-, C(NH)-NR<sub>8</sub>, C(=NR<sub>8</sub>)NR<sub>9</sub>, NR<sub>8</sub>-C(NH)-, NHR<sub>8</sub>(CO)NHR<sub>9</sub>, (O-CR<sub>8</sub>-CR<sub>8</sub>-O)-, CX(R<sub>8</sub>)<sub>2</sub>, CX<sub>2</sub>R<sub>8</sub>, CX<sub>3</sub>OCX<sub>3</sub>, N(R<sub>8</sub>)-S(O)R<sub>9</sub>, N(R<sub>8</sub>)-SO<sub>2</sub>R<sub>9</sub>, (=O), (=NH-OR<sub>8</sub>), C(=S)R<sub>8</sub>,NCR<sub>8</sub>, NOR<sub>8</sub>,NR<sub>8</sub>R<sub>9</sub>-SO<sub>2</sub>-NR<sub>8</sub>R<sub>9</sub>, OPO<sub>2</sub>R<sub>8</sub>, P*(R<sub>8</sub>)<sub>3</sub>X,SO<sub>3</sub>H, fused heteroaryl, and (=CR<sub>8</sub>) wherein R<sub>8</sub> and R<sub>9</sub> are independently H or C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, aralkyl, alkylaryl, cycloalkyl or heterocyclic, and X is a halogen which have excellent histamine-H<sub>3</sub> receptor antagonist activity as well as methods for preparing such compounds. In another embodiment, the invention discloses pharmaceutical compositions comprising such cyclic amines as well as methods of using them to treat allergy, inflammatory and CNS-related diseases.</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(a)</p> </div> <div style="text-align: center;">  <p>(b)</p> </div> <div style="text-align: center;">  <p>(c)</p> </div> <div style="text-align: center;">  <p>(d)</p> </div> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## N-(IMIDAZOLYLALKYL)SUBSTITUTED CYCLIC AMINES AS HISTAMINE-H<sub>3</sub> AGONISTS OR ANTAGONISTS

5 FIELD OF THE INVENTION


The present invention relates to N-(imidazolylalkyl) substituted cyclic amine compounds having valuable pharmacological properties, especially central nervous system ("CNS") activities and activity against inflammatory disease and allergic conditions. Compounds of this invention are agonists or antagonists of the histamine-H<sub>3</sub> receptor.

## BACKGROUND OF THE INVENTION

H<sub>3</sub> receptor sites are known and are of current interest to those skilled in the art as a therapeutic target. U.S. Patent 4,767,778 (Arrang *et al*) discloses certain imidazoles that behave as agonists of the H<sub>3</sub> receptors in rat brain. European Patent Application No. 0 420 396 A2 (Smith Kline & French Laboratories Limited) and Howson *et al.*, (*Bioorg. & Med. Chem. Letters*, (1992), Vol. 2 No. 1, pp. 77-78) describe imidazole derivatives having an amidine group as H<sub>3</sub> agonists. Van der Groot *et al.* (*Eur. J. Med. Chem.* (1992) Vol. 27, pp. 511-517) describe isothioureia analogs of histamine as potent agonists or antagonists of the histamine-H<sub>3</sub> receptor, and these isothioureia analogs of histamine overlap in part with those of the two references cited above. Clapham *et al.* ["Ability of Histamine-H<sub>3</sub> Receptor Antagonists to Improve Cognition and to Increase Acetylcholine Release *in vivo* in the Rat", *British Assn. for Psychopharmacology*, July 25-28 (1993), reported in *J. Psychopharmacol.* (Abstr. Book), A17] describe the ability of histamine-H<sub>3</sub> receptor antagonists to improve cognition and to increase release of acetylcholine *in vivo* in the rat. Clapham *et al.* ["Ability of the selective Histamine-H<sub>3</sub> Receptor Antagonist Thioperamide to improve Short-term Memory and Reversal Learning in the Rat", *Brit. J. Pharm. Suppl.*, 1993, 110, Abstract 65P] present results showing that thioperamide can improve short-term memory and reversal learning in the rat and implicate the involvement of H<sub>3</sub> receptors in the modulation of cognitive function. Yokoyama *et al.* ["Effect of Thioperamide, a Histamine-H<sub>3</sub> Receptor Antagonist, on Electrically Induced Convulsions in Mice", *Eur. J. Pharmacol.*, (1993), Vol. 234, pp. 129-133] report how thioperamide decreased the duration of each phase of convulsion and raised the electroconvulsive threshold, and

go on to suggest that these and other findings support the hypothesis that the central histaminergic system is involved in the inhibition of seizures. International Patent Publication No. WO 9301812-A1 (SmithKline Beecham PLC) describes the use of S-[3-(4(5)-imidazolyl)propyl]isothioureia as a histamine-H<sub>3</sub> antagonist, especially for treating cognitive disorders, *e.g.* Alzheimer's disease and age-related memory impairment. Schlicker *et al.* ["Novel Histamine-H<sub>3</sub> Receptor Antagonists: Affinities in an H<sub>3</sub> Receptor Binding Assay and Potencies in Two Functional H<sub>3</sub> Receptor Models", *British J. Pharmacol.*, (1994), Vol. 112, 1043-1048] describe a number of imidazolylalkyl compounds wherein the imidazolylalkyl group is bonded to a guanidine group, an ester group, an amide group, a thioamide group and a urea group, and compared these to thioperamide. Leurs *et al.* ["The Histamine-H<sub>3</sub>-receptor: A Target for Developing New Drugs", *Progr. Drug Res.* (1992), Vol. 39, pp. 127-165] and Lipp *et al.* ["Pharmacochemistry of H<sub>3</sub>-receptors" in *The Histamine Receptor*, eds.: Schwartz and Haas, Wiley-Liss, New York (1992), pp. 57-72] review a variety of synthetic H<sub>3</sub> receptor antagonists, and Lipp *et al.* (*ibid.*) have proposed the necessary structural requirements for an H<sub>3</sub> receptor antagonist.

20 WO 95/14007 claims H<sub>3</sub> receptor antagonists of the formula




wherein A, m, n, R<sup>1</sup> and R<sup>2</sup> are defined therein. The compounds are disclosed as being useful for treating various disorders, in particular such caused by allergy-induced responses.

25 WO 93/12093 discloses imidazolylmethyl piperazines and  
diazepines as H<sub>3</sub> antagonists. U.S. patent application, Serial No.  
08/965,754, filed November 7, 1997, discloses imidazolylalkyl substituted  
heterocyclic ring compounds as H<sub>3</sub> receptor antagonists. U.S. patent  
application, Serial No. 08/966,344, filed November 7, 1997, discloses  
30 phenylalkylimidazoles as H<sub>3</sub> receptor antagonists.

Reference is also made to US Application, Serial No. 08/689,951, filed August 16, 1996 which claims the combined use of a histamine-H<sub>1</sub> receptor antagonist and a histamine-H<sub>3</sub> receptor antagonist for treatment of allergy-induced airway responses.

- 3 -

Reference is also made to J. R. Bagley *et al*, *Journal of Medicinal Chemistry*, (1991), Vol. 34, 827-841, which discloses, among others, N-(imidazolylalkyl) substituted cyclic amine compounds useful as analgesics such as the amine compound with the formula:



5


In view of the art's interest in compounds which affect the  $H_3$  receptors, novel compounds having agonist or antagonist activity on  $H_3$  receptors would be a welcome contribution to the art. This invention provides just such a contribution by providing novel compounds having  $H_3$  agonist or antagonist activity.

10

### SUMMARY OF THE INVENTION

In one embodiment, this invention provides novel imidazole compounds having  $H_3$  agonist or antagonist activity. The inventive compounds are N-(imidazolylalkyl)-substituted cyclic amines having the general Formula I:

15



Formula I

wherein  $R_1$  is selected from the group consisting of H,  $-OCO-R_7$ , and  $-CO_2R_7$ , with  $R_7$  being a substituted or unsubstituted alkyl or aryl, the substituents on said substituted alkyl or aryl being selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic and halogen;

20




$R_2$  is selected from the group consisting of H, hydroxyl or halogen;  
 $G$  is a spacer moiety selected from the group consisting of a  $C_3-C_7$  straight chain alkyl, a  $C_2-C_7$  alkene or a  $C_2-C_7$  alkyne, with said straight

25


- 4 -

chain alkyl, alkene or alkyne being optionally substituted with one or more  $R_7$  groups; and

T which represents a cyclic amine is a ring moiety selected from the group consisting of a, b, c or d below:

abc

and

d

wherein n is an integer from 0-3, and  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be the same or different with the proviso that any two of said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be bound to the same or different carbon atom of ring T with a maximum valency of four at any carbon, with said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  being independently selected from the group consisting of H, substituted or unsubstituted  $C_1$ - $C_6$  alkyl, substituted or unsubstituted  $C_1$ - $C_6$  alkenyl, substituted or unsubstituted  $C_1$ - $C_6$  alkynyl, aryl, aralkyl, alkylaryl, cycloalkyl, heterocyclic,  $-C(O)R_8$ ,  $-C(=O)R_8$ ,  $-CO_2R_8$ ,  $-OC(O)R_8$ ,  $-NH_2$ ,  $-NHR_8$ ,  $-N(R_8)_2$ ,  $-NR_8R_9$ ,  $-SR_8$ ,  $-SO_2R_8$ ,  $S(O)R_8$ ,  $-OH$ ,  $-OR_8$ ,  $-CH_2OR_8$ ,  $-CH_2NH_2$ ,  $-CH_2N(R_8)_2$ ,  $-CH_2NHR_8$ ,  $-CH_2SR_8$ ,  $-C(O)NHR_8$ ,  $-C(O)NR_8R_9$ ,  $-CN$ ,  $-NO_2$ ,  $-NR_8-CO-$ ,  $-C(NH)-NR_8$ ,  $-C(=NR_8)NR_9$ ,  $-NR_8-C(NH)-$ ,  $-NHR_8(CO)NHR_9$ ,  $-(O-CR_8-CR_8-O)-$ ,  $-CX(R_8)_2$ ,  $-CX_2R_8$ ,  $-CX_3$ ,  $-OCX_3$ ,  $-N(R_8)-S(O)R_9$ ,  $-N(R_8)-SO_2R_9$ ,  $(=O)$ ,  $(=NH-OR_8)$ ,  $-C(=S)R_8$ ,  $-NCR_8$ ,  $-NOR_8$ ,  $-NR_8R_9-SO_2-NR_8R_9$ ,  $-OPO_2R_8$ ,  $-P^+(R_8)_3 X^-$ ,  $-SO_3H$ , fused heteroaryl, and  $(=CR_8)$  wherein  $R_8$  and  $R_9$  are independently H or  $C_1$ - $C_6$  alkyl, aryl, aralkyl, alkylaryl, cycloalkyl or heterocyclic, and X is a halogen.

Also included in the invention are tautomers, enantiomers and other optical isomers of compounds of Formula I, as well as pharmaceutically acceptable salts and solvates thereof.

- 5 -

A further feature of the invention is pharmaceutical compositions containing as active ingredient a compound of Formula I (or salt, solvate or isomers) together with a pharmaceutically acceptable carrier or excipient.


- 5 The invention also provides methods for preparing compounds of Formula I, as well as methods for treating diseases such as, for example, inflammation, allergy, diseases of the GI-tract, cardiovascular disease, or disturbances of the central nervous system. The latter methods comprise administering to a patient suffering from said disease or diseases a  
 10 therapeutically effective amount of a compound of Formula I, or pharmaceutical compositions comprising a compound of Formula I.

The invention also includes the aspect of using the claimed cyclic amine compound in combination with a histamine-H<sub>1</sub> receptor antagonist for treatment of allergy-induced airway (e.g., upper airway) responses.

15

### DETAILED DESCRIPTION OF THE INVENTION

In one embodiment, the present invention provides novel N-imidazolylalkyl substituted cyclic amine compounds of Formula I:



20

Formula I

- wherein R<sub>1</sub> is selected from the group consisting of H, -OCO-R<sub>7</sub>, and -CO<sub>2</sub>R<sub>7</sub>, with R<sub>7</sub> being a substituted or unsubstituted alkyl or aryl, the substituents on said substituted alkyl or aryl being selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl,  
 25 heterocyclic and halogen;




R<sub>2</sub> is selected from the group consisting of H, hydroxyl or halogen;

- G is a spacer moiety selected from the group consisting of a C<sub>3</sub>-C<sub>7</sub> straight chain alkyl, a C<sub>2</sub>-C<sub>7</sub> alkene or a C<sub>2</sub>-C<sub>7</sub> alkyne, with said straight chain alkyl, alkene or alkyne being optionally substituted with one or more  
 30 R<sub>7</sub> groups; and


T which represents a cyclic amine is a ring moiety selected from the group consisting of a, b, c or d below:



- 6 -

abc

and

d


- wherein  $n$  is an integer from 0-3, and  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be the same or different with the proviso that any two of said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be bound to the same or different carbon atom of ring T, with said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  being independently selected from the group consisting of H, substituted or unsubstituted  $C_1$ - $C_6$  alkyl, substituted or unsubstituted  $C_1$ - $C_6$  alkenyl, substituted or unsubstituted  $C_1$ - $C_6$  alkynyl, aryl, aralkyl, alkylaryl, cycloalkyl, heterocyclic,  $-C-O-R_8$ ,  $-C(=O)R_8$ ,  $-CO_2R_8$ ,  $-OC(O)R_8$ ,  $-NH_2$ ,  $-NHR_8$ ,  $-N(R_8)_2$ ,  $-NR_8R_9$ ,  $-SR_8$ ,  $-SO_2R_8$ ,  $S(O)R_8$ ,  $-OH$ ,  $-OR_8$ ,  $-CH_2OR_8$ ,  $-CH_2NH_2$ ,  $-CH_2N(R_8)_2$ ,  $-CH_2NHR_8$ ,  $-CH_2SR_8$ ,  $-C(O)-NHR_8$ ,  $-C(O)NR_8R_9$ ,  $-CN$ ,  $-NO_2$ ,  $-NR_8-CO-$ ,  $-C(NH)-NR_8$ ,  $-C(=NR_8)NR_9$ ,  $-NR_8-C(NH)-$ ,  $-NHR_8(CO)NHR_9$ ,  $-(O-CR_8-CR_8-O)-$ ,  $-CX(R_8)_2$ ,  $-CX_2R_8$ ,  $-CX_3$ ,  $-OCX_3$ ,  $-N(R_8)-S(O)R_9$ ,  $-N(R_8)-SO_2R_9$ ,  $(=O)$ ,  $(=NH-OR_8)$ ,  $-C(=S)R_8$ ,  $-NCR_8$ ,  $-NOR_8$ ,  $-NR_8R_9-SO_2-NR_8R_9$ ,  $-OPO_2R_8$ ,  $-P^+(R_8)_3 X^-$ ,  $-SO_3H$ , fused heteroaryl, and  $(=CR_8)$  wherein  $R_8$  and  $R_9$  are independently H or  $C_1$ - $C_6$  alkyl, aryl, aralkyl, alkylaryl, cycloalkyl or heterocyclic, and  $X$  is a halogen.

- Preferred moieties for  $R_1$  are hydrogen or a readily hydrolyzable group such as, for example,  $-O-CO-R_7$ , where  $R_7$  is described above.
- Preferred moieties for  $R_2$  are hydrogen or halogen.



Preferred representatives of  $G$  are alkyl groups such as  $n$ -propyl,  $n$ -butyl as well as their isomers.

Preferred cyclic amine ring for T is a piperidine ring ( $n=1$  in Formula I), as well as its 3,4-dehydro derivative, both being shown below:




- 7 -



Preferred moieties for  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are independently hydrogen, lower alkyl,  $-\text{CO-NHR}_8$ ,  $-\text{COOR}_8$ ,  $-\text{CH}_2\text{NH}-$ ,  $-\text{CH}_2\text{OH}$ ,  $-\text{NH}-$ ,  $-\text{CO}-$ ,  $-\text{CH}_2\text{-NR}_9\text{-SO}_2-$ , as well as sets of two moieties that attach to the same carbon atom such as, for example, acetal. Representative compounds of the invention include:

Formula IIFormula III

- 8 -

Formula IVFormula VFormula VI

- The compounds of the invention are basic and form
- 5 pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for such salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are


- 9 -

prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their corresponding salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their corresponding free base forms for purposes of this invention.

Depending upon the substituents on the inventive compounds, one may be able to form salts with bases too. Thus, for example, if there are carboxylic acid substituents in the molecule, salts may be formed with inorganic as well as organic bases such as, for example, NaOH, KOH,  $\text{NH}_4\text{OH}$ , tetraalkylammonium hydroxide, and the like.

As stated earlier, the invention includes tautomers, enantiomers and other stereoisomers of the compounds also. Thus, as one skilled in the art knows, certain imidazoles may exist in tautomeric forms. Such variations are contemplated to be within the scope of the invention.


Another embodiment of the invention discloses a method of making the N-(imidazolylalkyl)-substituted cyclic amines disclosed above. The compounds may be prepared by several processes well known in the art. A preferred method may be exemplified by considering the imidazole-containing part of the molecule as the "left hand side" and the cyclic amine part as the "right hand side", the latter being illustrated, for convenience sake, as:



In a preferred process, the left hand side part is taken as an aldehyde compound which is condensed with the right hand side part to form a Schiffs base (an enamine). A catalyst is not usually necessary; however, if one is desired, materials such as, for example, triethylaluminum, molecular sieves (4Å size, for example) may be used. Generally, this reaction is conducted in an organic solvent such as, for example, halogenated hydrocarbons, toluene, acetonitrile and the like, preferably in the absence of moisture, at temperatures in the range about 0-100°C by bringing together the ingredients in contact in the solvent

- 10 -

medium and stirring for about 10 minutes to about 48 hours at such temperatures. If the starting imidazole contains moieties such as, for example, NH, OH, CO<sub>2</sub>H and the like, such moieties are generally protected prior to the condensation reaction by using protecting groups such as, for example, triphenylmethyl, tert-butoxy, tert-butylester and the like, as is well known to those skilled in the art. The resulting Schiff's base may be isolated and reduced, or preferably reduced *in situ*, to the amine by treating with a suitable reductant such as, for example, sodium borohydride, sodium triacetoxyborohydride, sodium cyanoborohydride and the like, in an organic solvent at about 0-100°C for about 0.1-48 hours. At a desired time, any protecting group may be deprotected. Thus, for example, if the protecting group is N-triphenylmethyl, deprotection may be accomplished by acid hydrolysis. The reaction steps may be depicted by the scheme:



In the scheme above, PG represents a protecting group, and X represents (G-1) number of spacer moieties.

Isolation of the compound at various stages of the reaction may be achieved by standard techniques such as, for example, filtration, evaporation of solvent and the like. Purification of the product, intermediate and the like, may also be performed by standard techniques such as recrystallization, distillation, sublimation, chromatography, conversion to a suitable derivative which may be recrystallized and converted back to the amine, and the like. Such techniques are well known to those skilled in the art.

The thus prepared compounds may be analyzed for their composition and purity as well as characterized by standard analytical techniques such as, for example, elemental analysis, NMR, mass spectroscopy, and IR spectra.

- 11 -

The inventive compounds can readily be evaluated to determine activity at  $H_3$  receptors by known methods, including, for example, the guinea pig brain membrane assay and the guinea pig neuronal ileum contraction assay, both of which are described in U.S. patent 5,352,707.

5 Another useful assay utilizes rat brain membranes and is described by West *et al*, ("Identification of Two  $H_3$ -Histamine Receptor Subtypes", *Molecular Pharmacology*, (1990), Vol. 33, 610-613. Several of the present compounds were found to have high  $H_3$  antagonist activity some of which are disclosed in the **EXAMPLES** section below.

10 In another embodiment, this invention provides pharmaceutical compositions comprising the above-described N-(imidazolylalkyl) substituted cyclic amines as an active ingredient. The pharmaceutical compositions generally additionally comprise a pharmaceutically acceptable carrier diluent, excipient or carrier (collectively referred to  
15 herein as carrier materials). Because of their  $H_3$  antagonist activity, such pharmaceutical compositions possess utility in treating allergy, inflammation, nasal congestion, hypertension, glaucoma, sleeping disorders, states of hyper- and hypo-motility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers,  
20 Schizophrenia, migraines and the like diseases.

In yet another embodiment, the present invention discloses methods for preparing pharmaceutical compositions comprising the N-(imidazolylalkyl) substituted cyclic amines as an active ingredient. In the pharmaceutical compositions and methods of the present invention, the  
25 active ingredients will typically be administered in admixture with suitable carrier materials suitably selected with respect to the intended form of administration, i.e. oral tablets, capsules (either solid-filled, semi-solid filled or liquid filled), powders for constitution, oral gels, elixirs, dispersible granules, syrups, suspensions, and the like, and consistent with  
30 conventional pharmaceutical practices. For example, for oral administration in the form of tablets or capsules, the active drug component may be combined with any oral non-toxic pharmaceutically acceptable inert carrier, such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol,  
35 ethyl alcohol (liquid forms) and the like. Moreover, when desired or needed, suitable binders, lubricants, disintegrating agents and coloring agents may also be incorporated in the mixture. Powders and tablets may be comprised of from about 5 to about 95 percent inventive composition.

- 12 -

Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethylcellulose, polyethylene glycol and waxes. Among the lubricants there may be mentioned for use in these dosage forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrants include starch, methylcellulose, guar gum and the like. Sweetening and flavoring agents and preservatives may also be included where appropriate. Some of the terms noted above, namely disintegrants, diluents, lubricants, binders and the like, are discussed in more detail below.

Additionally, the compositions of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects, i.e. antihistaminic activity and the like. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injections or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier such as inert compressed gas, e.g. nitrogen.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides such as cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein by stirring or similar mixing. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby solidify.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

- 13 -

The compounds of the invention may also be deliverable transdermally. The transdermal compositions may take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as a re conventional in the art for this purpose.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active components, e.g., an effective amount to achieve the desired purpose.

The quantity of the inventive active composition in a unit dose of preparation may be generally varied or adjusted from about 0.01 milligrams to about 1,000 milligrams, preferably from about 0.01 to about 950 milligrams, more preferably from about 0.01 to about 500 milligrams, and typically from about 1 to about 250 milligrams, according to the particular application. The actual dosage employed may be varied depending upon the patient's age, sex, weight and severity of the condition being treated. Such techniques are well known to those skilled in the art. Generally, the human oral dosage form containing the active ingredients can be administered 1 or 2 times per day. The amount and frequency of the administration will be regulated according to the judgment of the attending clinician. A generally recommended daily dosage regimen for oral administration may range from about 0.04 milligrams to about 4,000 milligrams per day, in single or divided doses.

Capsule - refers to a special container or enclosure made of methyl cellulose, polyvinyl alcohols, or denatured gelatins or starch for holding or containing compositions comprising the active ingredients. Hard shell capsules are typically made of blends of relatively high gel strength bone and pork skin gelatins. The capsule itself may contain small amounts of dyes, opaquing agents, plasticizers and preservatives.

Tablet- refers to a compressed or molded solid dosage form containing the active ingredients with suitable diluents. The tablet can be prepared by compression of mixtures or granulations obtained by wet granulation, dry granulation or by compaction.

Oral gels-refers to the active ingredients dispersed or solubilized in a hydrophilic semi-solid matrix.



- 14 -

Powders for constitution refers to powder blends containing the active ingredients and suitable diluents which can be suspended in water or juices.

Diluent - refers to substances that usually make up the major portion of the composition or dosage form. Suitable diluents include sugars such as lactose, sucrose, mannitol and sorbitol; starches derived from wheat, corn rice and potato; and celluloses such as microcrystalline cellulose. The amount of diluent in the composition can range from about 10 to about 90% by weight of the total composition, preferably from about 25 to about 75%, more preferably from about 30 to about 60% by weight, even more preferably from about 12 to about 60%.

Disintegrants - refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Suitable disintegrants include starches; "cold water soluble" modified starches such as sodium carboxymethyl starch; natural and synthetic gums such as locust bean, karaya, guar, tragacanth and agar; cellulose derivatives such as methylcellulose and sodium carboxymethylcellulose; microcrystalline celluloses and cross-linked microcrystalline celluloses such as sodium croscarmellose; alginates such as alginic acid and sodium alginate; clays such as bentonites; and effervescent mixtures. The amount of disintegrant in the composition can range from about 2 to about 15% by weight of the composition, more preferably from about 4 to about 10% by weight.

Binders - refers to substances that bind or "glue" powders together and make them cohesive by forming granules, thus serving as the "adhesive" in the formulation. Binders add cohesive strength already available in the diluent or bulking agent. Suitable binders include sugars such as sucrose; starches derived from wheat, corn rice and potato; natural gums such as acacia, gelatin and tragacanth; derivatives of seaweed such as alginic acid, sodium alginate and ammonium calcium alginate; cellulosic materials such as methylcellulose and sodium carboxymethylcellulose and hydroxypropylmethylcellulose; polyvinylpyrrolidone; and inorganics such as magnesium aluminum silicate. The amount of binder in the composition can range from about 2 to about 20% by weight of the composition, more preferably from about 3 to about 10% by weight, even more preferably from about 3 to about 6% by weight.

Lubricant - refers to a substance added to the dosage form to enable the tablet, granules, etc. after it has been compressed, to release

- 15 -

from the mold or die by reducing friction or wear. Suitable lubricants include metallic stearates such as magnesium stearate, calcium stearate or potassium stearate; stearic acid; high melting point waxes; and water soluble lubricants such as sodium chloride, sodium benzoate, sodium acetate, sodium oleate, polyethylene glycols and d'l-leucine. Lubricants are usually added at the very last step before compression, since they must be present on the surfaces of the granules and in between them and the parts of the tablet press. The amount of lubricant in the composition can range from about 0.2 to about 5% by weight of the composition, preferably from about 0.5 to about 2%, more preferably from about 0.3 to about 1.5% by weight.

Glidants - materials that prevent caking and improve the flow characteristics of granulations, so that flow is smooth and uniform. Suitable glidants include silicon dioxide and talc. The amount of glident in the composition can range from about 0.1% to about 5% by weight of the total composition, preferably from about 0.5 to about 2% by weight.

Coloring agents - excipients that provide coloration to the composition or the dosage form. Such excipients can include food grade dyes and food grade dyes adsorbed onto a suitable adsorbent such as clay or aluminum oxide. The amount of the coloring agent can vary from about 0.1 to about 5% by weight of the composition, preferably from about 0.1 to about 1%.

Bioavailability - refers to the rate and extent to which the active drug ingredient or therapeutic moiety is absorbed into the systemic circulation from an administered dosage form as compared to a standard or control.

Conventional methods for preparing tablets are known. Such methods include dry methods such as direct compression and compression of granulation produced by compaction, or wet methods or other special procedures. Conventional methods for making other forms for administration such as, for example, capsules, suppositories and the like are also well known.

Another embodiment of the invention discloses use of the pharmaceutical compositions disclosed above for treatment of diseases such as, for example, allergy, inflammation, nasal congestion, hypertension, glaucoma, sleeping disorders, states of hyper- and hypomotility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers, Schizophrenia, migraines and the like. The method comprises administering a therapeutically effective amount of the

- 16 -

inventive pharmaceutical composition to a patient having such a diseases or diseases and in need of such a treatment.

In a still another embodiment, this invention discloses pharmaceutical compositions comprising the inventive N-(imidazolylalkyl) substituted cyclic amine compound in combination with a histamine-H<sub>1</sub> receptor antagonist. Optionally there may be a suitable pharmaceutically acceptable carrier present. Numerous chemical substances are known to have histamine-H<sub>1</sub> receptor antagonist activity. Many such compounds can be classified broadly as ethanolamines, ethylenediamines, alkylamines, phenothiazines, piperidines, and the like. Illustrative H<sub>1</sub> receptor antagonists useful in the practice of the present invention include, without limitation, astemizole, azatadine, azelastine, acrivastine, brompheniramine, chlorpheniramine, clemastine, cyclizine, carebastine, cyproheptadine, carbinoxamine, descarboethoxyloratadine (also known as desloratadine or "DCL"), doxylamine, dimethindene, ebastine, epinastine, efletirizine, fexofenadine, hydroxyzine, ketotifen, loratadine, levocabastine, meclizine, mizolastine, mequitazine, mianserin, noberastine, norastemizole, picumast, pyrilamine, promethazine, terfenadine, tripelennamine, temelastine, trimeprazine, and tripolidine. Other compounds can readily be evaluated to determine activity at H<sub>1</sub> receptors by known methods including, for example, specific blockade of the contractile response to histamine of isolated guinea pig ileum. All such H<sub>1</sub> receptor antagonists are suitable to prepare the pharmaceutical compositions.

Yet another embodiment of the invention discloses methods for preparing pharmaceutical compositions comprising the N-(imidazolylalkyl) substituted cyclic amines and a histamine-H<sub>1</sub> receptor antagonist. And a still another embodiment discloses the aspect of using such compositions for treatment of allergy-induced airway (e.g. upper airway) responses.

Those skilled in the art will realize that the term "upper airway" means the upper respiratory system--i.e., the nose, throat, and associated structures.


It will be apparent to those skilled in the art that many modifications, variations and alterations to the present disclosure, both to materials and methods, may be practiced. Such modifications, variations and alterations are intended to be within the spirit and scope of the present invention.

The following **EXAMPLES** are being provided to further illustrate the present invention. They are for illustrative purposes only; the scope of the invention is not to be considered limited in any way thereby.

**EXAMPLES**

Unless otherwise stated, the following abbreviations have the stated meanings in the Examples below:

- 5 DBU= 1,8-diazabicyclo[5.4.0]undec-7-ene  
 DBN= 1,5-diazabicyclo[4.3.0]non-5-ene  
 EDCI= 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
 HOBT= 1-hydroxybenzotriazole  
 DCC= dicyclohexylcarbodiimide  
 10 Dibal-H= diisobutylaluminum hydride  
 LAH= lithium aluminum hydride  
 NaBH(OAc)<sub>3</sub>= sodium triacetoxyborohydride  
 NaBH<sub>4</sub>= sodium borohydride  
 NaBH<sub>3</sub>CN= sodium cyanoborohydride  
 15 LDA= lithium diisopropylamide  
 p-TsOH= p-toluenesulfonic acid  
 TMAD= N,N,N',N'-tetramethylazodicarboxamide  
 CSA= camphorsulfonic acid  
 HRMS= High Resolution Mass Spectrometry  
 20 HPLC= High Performance Liquid Chromatography  
 LRMS= Low Resolution Mass Spectrometry  
 nM= nanomolar  
 K<sub>i</sub>= Dissociation Constant of Enzyme-Inhibitor complex  
 pA<sub>2</sub>= -logEC<sub>50</sub>, as defined by J. Hey, *Eur.J.Pharmacol.*, (1995), Vol.  
 25 294, 329.  
 Ci/mmol= Curie/mmol ( a measure of specific activity)  
 Tr= Triphenylmethyl  
 Tris= Tris(hydroxymethyl)aminomethane  
Example 1. Preparation of a compound of Formula I with  
 30 R<sub>1</sub>=triphenylmethyl, R<sub>2</sub>=R<sub>4</sub>=R<sub>5</sub>=R<sub>6</sub>=H, G=4, n=1, and R<sub>3</sub>=CO<sub>2</sub>Et:  
(i) Preparation of 1-triphenylmethyl-4-(4-butylcarboxaldehyde)  
imidazole:



- 18 -

To a solution of commercially available 4-imidazolecarboxaldehyde (50g, 0.52 mol) and triethylamine (190 mL) in methylene chloride (500 mL) was added triphenylmethyl chloride (175 g, 63 mol) in portions over 1.5 h. After 24 h the reaction was filtered, and the filtrate concentrated to approximately 300 mL. The organic phase was washed with water, 1 M NaOH, saturated brine, dried over MgSO<sub>4</sub> and concentrated in vacuo to a semi-solid. Recrystallization from hexanes-ethyl acetate afforded pure with 1-trityl imidazolecarboxaldehyde (55 g).


A 1 liter flask was charged with 1-trityl imidazolecarboxaldehyde (10 g, 29.6 mmol), THF (dry, 500 ml) and 3-benzyloxypropyl triphenylphosphonium bromide (15.3 g, 31.1 mmol) and cooled to -10 °C. Potassium t-Butoxide (1 M in dioxane, 31.2 mL) was added dropwise over 10 min and the reaction allowed to warm slowly to room temperature. After 4 h the reaction was filtered through celite, partially concentrated, ethyl acetate added and the organic layer washed with water. The organic layer was dried and concentrated to afford crude product.

A 500 mL hydrogenation vessel was charged with the crude product above (~ 11.5 g), methanol (60 mL), 10% palladium on carbon (1.0 g) and was shaken under hydrogen gas (55 psi) for 20 h. The catalyst was removed by filtration through celite and the product purified by chromatography on silica using methanol / methylene chloride eluent to afford 7.2 g with 4-hydroxypropyl 1-trityl imidazole.

A 500 mL flask was charged with methylene chloride, DMSO ((6 g, 76 mmol) and cooled to -65 °C. Oxalyl chloride ((7.2 g, 57 mmol) was added in three portions over 10 min. After 10 min., 4-hydroxypropyl-1-trityl imidazole (7.2 g, 19 mmol) was added as a solution in methylene chloride. After 5 min triethyl amine (9.6 g, 95 mmol) was added and the reaction allowed to warm slowly and stir overnight. The reaction was quenched with 1M potassium hydrogen phosphate, washed with water and the organic layer concentrated to crude aldehyde (5.7 g).


- 19 -

(ii) Preparation of a compound of Formula I with  $R_1$ =triphenylmethyl,  $R_2=R_4=R_5=R_6$ =H,  $G=4$ ,  $n=1$ , and  $R_3$ =CO<sub>2</sub>Et:




- 5 To a 1,2-dichloroethane solution (15 ml) of 1-triphenylmethyl-4-(4-butylcarboxaldehyde) imidazole (1.66 mg, 4.37 mmol) was added ethyl isonipecotinate (0.84 ml, 5.45 mmol) followed by sodium triacetoxyborohydride (1.39 g, 6.56 mmol) at room temperature. The mixture was stirred for 24 h under nitrogen and then quenched with water.
- 10 Sodium hydroxide (5% aq.) was added and the layers were separated. The combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The crude product was chromatographed on silica gel eluting with 2% MeOH-CH<sub>2</sub>Cl<sub>2</sub> increasing gradually to 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>. The product was obtained (2.03 g, 89%) as a clear oil
- 15 which solidified on standing. LRMS (CI, M+H) = 522.

- 20 -

Example 2: Deprotection of the product from Example 1:

To a dioxane solution (3 ml) of the trityl protected imidazole (170 mg, 0.330 mmol) was added a 4M HCl-dioxane solution (0.65 ml) at room temperature. The mixture was heated to 80 °C for 4 h, at which time the hydrochloride salt precipitated out of solution as a gum covering the walls of the flask. The mixture was cooled to room temperature, and the solvent was decanted from the gummy residue. The residue was rinsed consecutively with Et<sub>2</sub>O, EtOAc and CH<sub>2</sub>Cl<sub>2</sub> and dried under vacuum to provide the desired material as a light brown gum. LRMS (CI, M+H) = 280.


Example 3: Preparation of compound according to the reaction:



- 5 Trifluoromethoxyaniline (0.0285 ml, 0.211 mmol) was mixed with  
anhydrous  $\text{CH}_2\text{Cl}_2$  (1 ml) in a dry flask. Trimethyl aluminum (0.315 ml,  
0.633 mmol) was added via syringe, and the reaction mixture was stirred  
at room temperature for 15 min. A  $\text{CH}_2\text{Cl}_2$  solution of ester (100 mg, 0.192  
10 mmol) was then added and stirring was continued for 48 h. The reaction  
was quenched carefully with  $\text{H}_2\text{O}$  and MeOH, filtered through celite, and  
concentrated. Chromatography on silica gel eluting with 100%  $\text{CH}_2\text{Cl}_2$   
increasing gradually to 10% MeOH- $\text{CH}_2\text{Cl}_2$  afforded 94 mg (75%) of the  
desired material. LRMS (CI, M+H) = 653. Analytical CHN for  
( $\text{C}_{39}\text{H}_{39}\text{N}_4\text{O}_2\text{F}_3$ ): C, 69.65; H, 6.17; N, 8.33: Found C, 69.55; H, 6.19; N, 8.25.




Example 4: Preparation of compound of Formula II according to the reaction:




5 The trityl compound (162 mg, 0.248 mmol) was taken as a dioxane solution (3 ml). 4M HCl-dioxane solution (0.5 ml) was added and reacted as above. The product was obtained as a light brown gum. LRMS (CI, M+H) = 411. HRMS calc. 411.2008; Found 411.2011.

- 23 -

Example 5:


5  $n$ -Propylamine (0.226 ml, 2.76 mmol) was mixed with anhydrous toluene (10 ml) in a dry flask. Trimethyl aluminum (4.2 ml, 8.3 mmol) was added via syringe, and the reaction mixture was stirred at room temperature for 15 min. A toluene solution of the ester (1.3 g, 2.51 mmol) was then added and stirring was continued for 24 h at 60 °C. The reaction was cooled to room temperature and quenched carefully with  $\text{H}_2\text{O}$  and MeOH, filtered through celite, and concentrated. Chromatography on silica  
10 gel eluting with 100%  $\text{CH}_2\text{Cl}_2$  increasing gradually to 10% MeOH- $\text{CH}_2\text{Cl}_2$  afforded 1.28 g (95%) of the desired product. LRMS ( $\text{Cl}^-$ ,  $\text{M}+\text{H}$ ) = 535. Analytical CHN for ( $\text{C}_{35}\text{H}_{42}\text{N}_4\text{O} \times 0.8 \text{ H}_2\text{O}$ ): C, 76.55; H, 8.00; N, 10.20: Found C, 76.76; H, 7.99; N, 10.21.

### Example 6: Reduction of the Amide to Amine:



To a mixture of LAH (274 mg, 7.2 mmol) in freshly distilled THF (12 ml) was added a THF solution of the amide (1.27 g, 2.38 mmol) at room temperature. The mixture was then heated to reflux for 3.5 hr., then cooled to room temperature and quenched with solid  $\text{Na}_2\text{SO}_4 \cdot 10 \text{H}_2\text{O}$ , and 2.5M NaOH (0.5 ml). The slurry was filtered through celite and the filtrate was concentrated and chromatographed on silica gel (30 g) eluting with 10% MeOH- $\text{CH}_2\text{Cl}_2$  increasing to 10% MeOH- $\text{CH}_2\text{Cl}_2$  containing 1%  $\text{NH}_4\text{OH}$ , afforded 904 mg (73%) of product. LRMS (CI, M+H) = 521. HRMS calc. 521.3644; found 521.3643.


Example 7:



To a dioxane solution (3 ml) of the trityl precursor (117 mg, 0.219 mmol) was added 4M HCl-dioxane solution (0.5 ml) and reacted as above


- 5 The deprotected amide was obtained as a light brown gum. LRMS (CI, M+H) =293. HRMS calc. 293.2341; Found 293.2339.

Example 8:



To a dioxane solution (3 ml) of the trityl precursor (270 mg, 0.519 mmol) was added 4M HCl-dioxane solution (1.5 ml) and reacted as above. The product was obtained as a light brown gum. LRMS (CI, M+H) = 279. HRMS calc. 279.2549; Found 279.2549.


Example 9:



10 To a  $\text{CH}_2\text{Cl}_2$  (3 ml) containing  $\text{Et}_3\text{N}$  (0.217 ml, 1.56 mmol) and the imidazolylamine (270 mg, 0.520 mmol) was added 4-

- 27 -


chlorobenzenesulfonyl chloride (108 mg, 0.510 mmol). The mixture was stirred at room temperature for 24 h, then concentrated, and chromatographed directly on silica gel (25 g) eluting with 5% MeOH-CH<sub>2</sub>Cl<sub>2</sub> increasing to 5% MeOH-CH<sub>2</sub>Cl<sub>2</sub> containing 1% NH<sub>4</sub>OH which  
5 afforded 272 mg (77%) of pure material. LRMS (CI, M+H) = 695. Analytical CHNS for (C<sub>41</sub>H<sub>45</sub>N<sub>4</sub>O<sub>2</sub>SCI x 0.7 H<sub>2</sub>O): C, 69.56; H, 6.89; N, 7.91; S, 4.53: Found C, 69.53; H, 7.09; N, 7.72; S, 4.36.

Example 10:


10

To a dioxane solution (3 ml) of the trityl precursor (192 mg, 0.277 mmol) was added 4M HCl-dioxane solution (0.6 ml) and reacted as above. The product was obtained as a light brown gum. LRMS (CI, M+H) = 453.

- 28 -

Example 11:


- To a  $\text{CH}_2\text{Cl}_2$  (3 ml) containing  $\text{Et}_3\text{N}$  (0.217 ml, 1.56 mmol) and the amine (270 mg, 0.520 mmol) was added 4-trifluoromethoxybenzene sulfonyl chloride (0.088 ml, 0.510 mmol). The mixture was stirred at room temperature for 24 h, then concentrated, and chromatographed directly on silica gel (25 g) eluting with 5%  $\text{MeOH}-\text{CH}_2\text{Cl}_2$  increasing to 5%  $\text{MeOH}-\text{CH}_2\text{Cl}_2$  containing 1%  $\text{NH}_4\text{OH}$  which afforded 171 mg (45%) of pure material. LRMS (CI,  $\text{M}+\text{H}$ ) = 745. HRMS calc for  $(\text{C}_{42}\text{H}_{47}\text{N}_4\text{O}_3\text{SF}_3)$  745.3399; Found 745.3417.

Example 12: Preparation of compound of Formula IV:

- 29 -

To a dioxane (3 ml) solution of the trityl precursor (136 mg, 0.183 mmol) was added 4M HCl-dioxane solution (0.6 ml) and reacted as above. The product was obtained as a light brown gum. LRMS (CI, M+H) = 503.

5


Example 13:

To the starting ester (500 mg, 0.959 mmol) were added p-chloroaniline (145 mg, 1.13 mmol), toluene (5 ml), trimethylaluminum (2.0 ml, 4.0 mmol) and reacted as above. The product was obtained (486 mg, 83%) as a faint yellow solid. LRMS (CI, M+H) = 603.

10




- 30 -

Example 14:

To the ester (500 mg, 0.959 mmol) were added 3,4-dichloroaniline (190 mg, 1.17 mmol), toluene (5 ml), trimethylaluminum (2.0 ml, 4.0 mmol) and reacted as above. The product was obtained (530 mg, 87%) as a faint amber solid. LRMS (Cl, M+H) = 637.


- 31 -

Example 15:




To the ester (500 mg, 0.959 mmol) were added m-dichloroaniline (150 mg, 1.18 mmol), toluene (5 ml), trimethylaluminum (2.0 ml, 4.0 mmol) and reacted as above. The product was obtained (557 mg, 96%) as an off white foam. LRMS (CI, M+H) = 603.

Example 16:




To the ester (500 mg, 0.959 mmol) were added 3-trifluoromethylaniline (185 mg, 1.15 mmol), toluene (5 ml),  
5 trimethylaluminum (2.0 ml, 4.0 mmol) and reacted as above. The product was obtained (533 mg, 88%) as an off white solid. MP = 145-148 °C. LRMS (CI, M+H) = 637.

Example 17:




To a dioxane solution (6 ml) of the trityl precursor (160 mg, 0.266 mmol) was added a 4M HCl-dioxane solution (0.5 ml) at room temperature. The mixture was heated to 80 °C for 4 h, at which time the hydrochloride salt precipitated out of solution as a gum covering the walls of the flask. The mixture was cooled to room temperature, and the solvent was decanted from the gummy residue. The residue was rinsed consecutively with Et<sub>2</sub>O, EtOAc and CH<sub>2</sub>Cl<sub>2</sub> and dried under vacuum to provide 92 mg of the product as a light brown gum. LRMS (Cl, M+H) = 361.

Example 18:



Using the same procedure as in Example 17, the trityl precursor (200 mg, 0.314 mmol) in dioxane (6 ml) was deprotected using 4M HCl-dioxane 0.5 ml). The product was obtained (138 mg) as a brown gum. LRMS (CI, M+H) = 395.


Example 19: Preparation of compound of Formula VI:



By using a similar procedure as in Example 17, the trityl precursor (200 mg, 0.332 mmol) in dioxane (6 ml) was deprotected using 4M HCl-dioxane (0.5 ml) to yield 126 mg of product as an amber foam. LRMS (Cl, M+H) = 361.

5

Example 20: Preparation of compound of Formula V:



By using a similar procedure as in Example 17, the trityl precursor (200 mg, 0.332 mmol), in dioxane (6 ml) was deprotected using 4M HCl-dioxane (1.0 ml) to yield 183 mg of product as an amber foam. LRMS (CI, M+H) = 395.

### General Procedure for H<sub>3</sub>-Receptor Binding Assay

15 The source of the H<sub>3</sub> receptors in this experiment was guinea pig  
brain. The animals weighed 400-600 g. The brain tissue was  
homogenized with a solution of 50 mM Tris, pH 7.5. The final  
concentration of tissue in the homogenization buffer was 10% w/v. The  
homogenates were centrifuged at 1,000 x g for 10 min. in order to remove  
clumps of tissue and debris. The resulting supernatants were then  
20 centrifuged at 50,000 x g for 20 min. in order to sediment the membranes,  
which were next washed three times in homogenization buffer (50,000 x g  
for 20 min. each). The membranes were frozen and stored at -70°C until  
needed.

- 36 -

All compounds to be tested were dissolved in DMSO and then diluted into the binding buffer (50 mM Tris, pH 7.5) such that the final concentration was 2  $\mu\text{g/ml}$  with 0.1% DMSO. Membranes were then added (400  $\mu\text{g}$  of protein) to the reaction tubes. The reaction was started by the addition of 3 nM [ $^3\text{H}$ ]R- $\alpha$ -methylhistamine (8.8 Ci/mmol) or 3 nM [ $^3\text{H}$ ]N $^{\alpha}$ -methylhistamine (80 Ci/mmol) and continued under incubation at 30°C for 30 min. Bound ligand was separated from unbound ligand by filtration, and the amount of radioactive ligand bound to the membranes was quantitated by liquid scintillation spectrometry. All incubations were performed in duplicate and the standard error was always less than 10%. Compounds that inhibited more than 70% of the specific binding of radioactive ligand to the receptor were serially diluted to determine a  $K_i$  (nM). The results are given in the Table 1 for the HCl salt of the indicated compound.

		<u>Table 1</u>				
	<u>No.</u>	<u>G</u>	<u>I</u>	<u>R</u>	<u>K<sub>i</sub></u>	<u>pA<sub>2</sub></u>
15						
	1	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	-COOEt	22	
20	2	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CONH-Ph-4-OCF <sub>3</sub> (Example 4)	2	9.3
	3	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CO-NH-Propyl (Example 7)	53	
25	4	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CH <sub>2</sub> -NH-Propyl (Example 8)	200	
	5	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CH <sub>2</sub> -NH- (SO <sub>2</sub> -Ph-4-Cl)-propyl (Example 10)	44	
30						
	6	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CH <sub>2</sub> -NH- (SO <sub>2</sub> -Ph-4-OCF <sub>3</sub> )-propyl (Example 12)	24	6.5
35						
	7	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CONH-Ph-4-Cl (Example 17)	35	
40						
	8	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CONH-Ph-3,4-Cl (Example 18)	27	6.8
	9	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CONH-Ph-3-Cl (Example 19)	5.5	8.9
45						
	10	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CONH-Ph-3-CF <sub>3</sub>	9	9.1

- 37 -

	11	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-Ph-3-OCF <sub>3</sub>	1	9.2
5	12	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-CH <sub>2</sub> Ph-4-Cl	29	8
	13	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-CH <sub>2</sub> Ph-4-OCF <sub>3</sub>	24	7.5
	14	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-Ph-4-OBu	10	9
10	15	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-Ph-4-OH	100	
	16	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CONH-Ph-3-OH	700	
15	17	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-NHCO-Ph-4-OCF <sub>3</sub>	23	
	18	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-NHCO-Ph-4-F	52	
	19	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CH <sub>3</sub> ; 4-CONH-Ph-4-OCF <sub>3</sub>	180	7.4
20	20	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	3-CH <sub>3</sub> ; 3-CONH-Ph-4-OCF <sub>3</sub>	170	
25	21	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-NH-CH <sub>2</sub> Ph-4-OCF <sub>3</sub>	13	8.1
	22	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CH <sub>2</sub> NH-Ph-4-OCF <sub>3</sub>	9	
	23	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-NH-Ph-3-OCF <sub>3</sub>	7	8.1
30	24	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-N-piperidine	100	6.2
	25	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-OH; 4-Ph-4-Cl	6	7.9
35	26	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-OH; 4-Ph-4-Cl	6	8
	27	(CH <sub>2</sub> ) <sub>5</sub>	<u>a</u>	4-OH; 4-Ph-4-Cl	5	8.1
	28	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-OH; 4-Ph-3-CF <sub>3</sub>	60	
40	29	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-(CH <sub>2</sub> ) <sub>3</sub> -4-piperidiny- (CH <sub>2</sub> ) <sub>2</sub> -OH	12	7.1
45	30	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-(CH <sub>2</sub> ) <sub>3</sub> -4-piperidiny- (CH <sub>2</sub> ) <sub>2</sub> -OH	7	7.7
	31	(CH <sub>2</sub> ) <sub>4</sub> (3,4-dihydro)	<u>c</u>	4-Ph-4-F	14	7.9
50	32	(CH <sub>2</sub> ) <sub>3</sub>	<u>a</u>	4-CO-Ph-4-F	6	7.7
	33	(CH <sub>2</sub> ) <sub>4</sub>	<u>a</u>	4-CO-Ph-4-F	3	8.2



- 38 -

	34	(CH <sub>2</sub> ) <sub>4</sub>	a	4-CH <sub>2</sub> -OPh-4-F	10	7.7
	35	(CH <sub>2</sub> ) <sub>3</sub>	a	4-Ph, 4-COEt	500	
5	36	(CH <sub>2</sub> ) <sub>3</sub>	a	4-OH, 4-CH <sub>2</sub> Ph	660	

From these test results and the background knowledge about the compounds described in the references in the section "Background of the Invention", it would be apparent to the skilled artisan that the compounds of the invention would be useful in treating inflammation, allergy, diseases of the GI-tract, cardiovascular disease, disturbances of the central nervous system, and the like diseases stated earlier.

The following Example A and Example B illustrate, in a non-limiting way, the preparation of pharmaceutical dosage forms which contain a compound of the invention. As used therein, the term "active compound" is used to designate one of the compounds of Formula I or salt thereof.

### Pharmaceutical Dosage Form Examples

#### EXAMPLE A

##### Tablets

No.	Ingredients	mg/tablet	mg/tablet
1	Active compound	100	500
2	Lactose USP	122	113
3	Corn Starch, Food Grade, as a 10% paste in Purified Water	30	40
4	Corn Starch, Food Grade	45	40
5	Magnesium Stearate	3	7
	Total	300	700

- 39 -

General Method of Manufacture

Items No. 1 and 2 are mixed in a suitable mixer for about 10 to 15 minutes. The mixture is then granulated with Item No. 3. The damp granules are milled through a coarse screen (e.g., 1/4", 0.63 cm) if necessary. The damp granules are then dried. The dried granules are screened if necessary and then mixed with Item No. 4 for about 10–15 minutes. Item No. 5 is then added and mixed for about 1 to 3 minutes. The mixture is then compressed to the appropriate size and weight on a suitable tablet machine.

EXAMPLE BCapsules

	<u>No.</u>	<u>Ingredient</u>	<u>mg/capsule</u>	<u>mg/capsule</u>
15	1	Active compound	100	500
	2	Lactose USP	106	123
	3	Corn Starch, food grade	40	70
20	4	Magnesium stearate, NF	4	7
	Total:		250	700

General Method of Manufacture


Items No. 1, 2 and 3 are mixed in a suitable blender for about 10 to 15 minutes. Item No. 4 is then added and mixed for about 1 to 3 minutes. The mixture is filled into suitable two-piece hard gelatin capsules on a suitable encapsulating machine.

- 40 -

**CLAIMS**

What is claimed is:

1. A compound, including enantiomers, stereoisomers and tautomers thereof, or pharmaceutically acceptable salts or solvates of said compound, said compound having the formula:






- wherein  $R_1$  is selected from the group consisting of H,  $-OCO-R_7$ , and  $-CO_2R_7$ , with  $R_7$  being a substituted or unsubstituted alkyl or aryl, the substituents on said substituted alkyl or aryl being selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic and halogen;


$R_2$  is selected from the group consisting of H, hydroxyl or halogen;

- G is a spacer moiety selected from the group consisting of a  $C_3$ - $C_7$  straight chain alkyl, a  $C_2$ - $C_7$  alkene or a  $C_2$ - $C_7$  alkyne, with said straight chain alkyl, alkene or alkyne being optionally substituted with one or more  $R_7$  groups; and

T which represents a cyclic amine is a ring moiety selected from the group consisting of a, b, c or d below:

abc


and

d

- 41 -


- wherein n is an integer from 0-3, and  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be the same or different with the proviso that any two of said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  can be bound to the same or different carbon atom of ring T, with said  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  being independently selected from the group consisting of H, substituted or unsubstituted  $C_1$ - $C_6$  alkyl, substituted or unsubstituted  $C_1$ - $C_6$  alkenyl, substituted or unsubstituted  $C_1$ - $C_6$  alkynyl, aryl, aralkyl, alkylaryl, cycloalkyl, heterocyclic,  $-C-O-R_8$ ,  $-C(=O)R_8$ ,  $-CO_2R_8$ ,  $-OC(O)R_8$ ,  $-NH_2$ ,  $-NHR_8$ ,  $-N(R_8)_2$ ,  $-NR_8R_9$ ,  $-SR_8$ ,  $-SO_2R_8$ ,  $S(O)R_8$ ,  $-OH$ ,  $-OR_8$ ,  $-CH_2OR_8$ ,  $-CH_2NH_2$ ,  $-CH_2N(R_8)_2$ ,  $-CH_2NHR_8$ ,  $-CH_2SR_8$ ,  $-C(O)-NHR_8$ ,  $-C(O)NR_8R_9$ ,  $-CN$ ,  $-NO_2$ ,  $-NR_8-CO-$ ,  $-C(NH)-NR_8$ ,  $-C(=NR_8)NR_9$ ,  $-NR_8-C(NH)-$ ,  $-NHR_8(CO)NHR_9$ ,  $-(O-CR_8-CR_8-O)-$ ,  $-CX(R_8)_2$ ,  $-CX_2R_8$ ,  $-CX_3$ ,  $-OCX_3$ ,  $-N(R_8)-S(O)R_9$ ,  $-N(R_8)-SO_2R_9$ ,  $(=O)$ ,  $(=NH-OR_8)$ ,  $-C(=S)R_8$ ,  $-NCR_8$ ,  $-NOR_8$ ,  $-NR_8R_9-SO_2-NR_8R_9$ ,  $-OPO_2R_8$ ,  $-P^+(R_8)_3 X^-$ ,  $-SO_3H$ , fused heteroaryl, and  $(=CR_8)$  wherein  $R_8$  and  $R_9$  are independently H or  $C_1$ - $C_6$  alkyl, aryl, aralkyl, alkylaryl, cycloalkyl or heterocyclic, and X is a halogen.

2. The compound of claim 1, wherein  $R_1$  is H, substituted or unsubstituted alkyl,  $-O-CO-R_7$  or  $-CO_2R_7$ .
3. The compound of claim 2, wherein  $R_1$  is H.
4. The compound of claim 2, wherein  $R_1$  is  $-CO_2R_7$ .
5. The compound of claim 1, wherein G is n-propyl.
6. The compound of claim 1, wherein G is n-butyl.
7. The compound of claim 1, wherein  $R_2$  is halogen.
8. The compound of claim 7, wherein  $R_2$  is F.
9. The compound of claim 1, wherein n is 1.
10. The compound of claim 1, wherein T is



wherein  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are as defined in claim 1.


- 11 The compound of claim 1, wherein T is



- 42 -

wherein  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are as defined in claim 1.

12. The compound of claim 1, wherein T is



wherein  $R_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  are as defined in claim 1.

5 13. The compound of claim 1, wherein  $R_3=R_4=H$  and  $R_5=O$ .

14. The compound of claim 1, wherein  $R_3$  and  $R_4$  are on the same carbon of ring T and are independently  $-CH_3$  and  $-C(O)-NH$ -aryl,  $R_5=O$  and  $R_6$  is H.

10 15. The compound of claim 1, wherein  $R_3$  is methyl,  $R_4$  is an amide and  $R_5=R_6=H$ .

16. The compound of claim 15, wherein said  $R_3$  and  $R_4$  are on the same carbon of ring T.


17. The compound of claim 1, wherein  $R_3$  and  $R_4$  are on the same carbon of ring T and are collectively  $-(O-CH_2-CH_2-O-)$ ,  $R_5=R_6=H$ .

15 18. The compound of claim 1, wherein  $R_3=CH_2OR_{10}$  and  $R_4=R_5=R_6=H$ , where  $R_{10}$  is H or  $CH_3$ .

19. The compound of claim 1, wherein  $R_3$  and  $R_4$  are on the same carbon of ring T and are independently  $-OH$  and 2-pyridyl, and  $R_5=R_6=H$ .


20 20. The compound of claim 1, wherein said pharmaceutically acceptable salt is a hydrochloride.

21. A compound of the formula:




22. A compound of the formula:

- 43 -



23. A pharmaceutical composition for use in treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper- and hypo-motility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers, Schizophrenia and migraines, said composition comprising a therapeutically effective amount of a compound of Claim 1 and a pharmaceutically acceptable carrier.
24. A method of treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper- and hypo-motility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers, Schizophrenia and migraines, said method comprising administering a therapeutically effective amount of a compound of claim 1 to a patient in need of such a treatment.
25. The use of a compound of claim 1 for the manufacture of a medicament for use in treating allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper- and hypo-motility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers, Schizophrenia and migraines.
26. The use of a compound of claim 1 for the treatment of allergy, inflammation, hypertension, glaucoma, sleeping disorders, states of hyper- and hypo-motility of the gastrointestinal tract, hypo- and hyper-activity of the central nervous system, Alzheimers, Schizophrenia and migraines.
27. A method of preparing a pharmaceutical composition comprising admixing a compound of claim 1 with a pharmaceutically acceptable carrier.
28. A method of preparing a compound of claim 1, as outlined in the scheme:

- 44 -



said method comprising: (i) reacting an imidazolylalkyl-substituted aldehyde with an amine of ring T in an organic solvent at a temperature of about -20°C to about 150°C to form an enamine, and (ii) reducing said enamine to yield an imidazolylalkyl substituted amine (of ring T), wherein R<sub>1</sub>, R<sub>2</sub> and T are as defined in claim 1.

29. The method of claim 28, wherein R<sub>1</sub> is H or triphenylmethyl, R<sub>2</sub>=H and ring T carries a -CO<sub>2</sub>Et substituent on position 4 with respect to said amine nitrogen.

30. The method of claim 29, wherein said triphenylmethyl is subsequently hydrolyzed to form R<sub>1</sub>=H.

31. The method of claim 29, further comprising reacting said -CO<sub>2</sub>Et substituent on ring T with an amine to form an amide.

32. The method of claim 31, further comprising reducing said amide to form a second amine substituent on said ring T.

33. The method of claim 32, further comprising reacting said second amine substituent with a sulfonyl halide to form a sulfonamide.

34. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 in admixture with or in combination with a therapeutically effective amount of a histamine-H<sub>1</sub> receptor antagonist.

35. The use of a compound of claim 1 in combination with or in admixture with a histamine-H<sub>1</sub> receptor antagonist for treatment of upper airway allergic responses.

36. A method for treatment of upper airway allergic responses comprising administering a compound of claim 1 in admixture with or in combination with a histamine-H<sub>1</sub> receptor antagonist.

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/21461

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 C07D401/06 C07D233/56 A61K31/4164 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STARK H ET AL: "Developments of Histamine H3-receptor Antagonists" DRUGS OF THE FUTURE, ES, BARCELONA, vol. 21, no. 5, 1 January 1996 (1996-01-01), pages 507-520, XP002084872 ISSN: 0377-8282 * see page 512, fig. 7 * the whole document	1-36
X	J.-C. SCHWARTZ, H.L. HAAS (EDS.): "The Histamine Receptor" 1992, WILEY-LISS, INC., NEW YORK XP002132481 cited in the application * see page 64, fig. 2 * page 57 -page 72	1-36

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"A" document member of the same patent family

Date of the actual completion of the international search

9 March 2000

Date of mailing of the international search report

27/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fac. (+31-70) 340-3016

Authorized officer

Stellmach, J



# INTERNATIONAL SEARCH REPORT

Intern. Patent Application No  
PCT/US 99/21461

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 40126 A (GLIATECH INC ; PHILLIPS JAMES G (US); KHAN AMIN MOHAMMED (US); TEDF) 19 December 1996 (1996-12-19) the whole document	1-36
Y	WO 93 12093 A (SCHERING CORP) 24 June 1993 (1993-06-24) the whole document	1-36
Y	STARK, H. ET AL.: "Novel Carbamates as Potent Histamine H3 Receptor Antagonists with High in vitro and Oral in vivo Activity" J.MED.CHEM., vol. 39, no. 5, 1996, pages 1157-1163, XP000882765 WASHINGTON * see page 1160, table 1, ex. 42 * the whole document	1-36
Y	GANELLIN, C.R. ET AL.: "A Novel Series of (Phenoxyalkyl)imidazoles as Potent H3-Receptor Histamine Antagonists" J.MED.CHEM., vol. 39, no. 19, 1996, pages 3806-3813, XP000882764 WASHINGTON the whole document	1-36
Y	EP 0 420 396 A (SMITH KLINE FRENCH LAB) 3 April 1991 (1991-04-03) the whole document	1-36
Y	WO 93 12107 A (SCHERING CORP) 24 June 1993 (1993-06-24) the whole document	1-36
Y	WO 93 12108 A (SCHERING CORP) 24 June 1993 (1993-06-24) the whole document	1-36

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/21461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640126 A	19-12-1996	AU 713166 B AU 6248296 A CA 2222652 A EP 0841923 A JP 11507631 T	25-11-1999 30-12-1996 19-12-1996 20-05-1998 06-07-1999
WO 9312093 A	24-06-1993	AT 128708 T AU 662588 B AU 3247793 A CA 2125964 A DE 69205287 D DE 69205287 T DK 618905 T EP 0618905 A ES 2078118 T GR 3018474 T HK 185696 A IL 104126 A JP 2540015 B JP 6511251 T KR 136207 B NZ 246295 A US 5633250 A ZA 9209786 A	15-10-1995 07-09-1995 19-07-1993 24-06-1993 09-11-1995 14-03-1996 04-12-1995 12-10-1994 01-12-1995 31-03-1996 11-10-1996 09-05-1999 02-10-1996 15-12-1994 25-04-1998 25-06-1996 27-05-1997 18-06-1993
EP 0420396 A	03-04-1991	DE 69003213 D DE 69003213 T JP 3058924 A US 5047418 A	14-10-1993 05-01-1994 14-03-1991 10-09-1991
WO 9312107 A	24-06-1993	AT 140223 T AU 665604 B AU 3275893 A CA 2126086 A DE 69212164 D DE 69212164 T DK 619818 T EP 0619818 A ES 2089782 T GR 3021195 T HK 31097 A IL 104124 A JP 7121938 B JP 6511252 T KR 142801 B MX 9207392 A NZ 246434 A SG 44565 A US 5932596 A ZA 9209785 A	15-07-1996 11-01-1996 19-07-1993 24-06-1993 14-08-1996 05-12-1996 25-11-1996 19-10-1994 01-10-1996 31-12-1996 21-03-1997 06-12-1998 25-12-1995 15-12-1994 15-07-1998 01-06-1993 27-02-1996 19-12-1997 03-08-1999 21-06-1993
WO 9312108 A	24-06-1993	AT 141269 T AU 671092 B AU 3277393 A CA 2129679 A DE 69212844 D DE 69212844 T DK 618913 T	15-08-1996 15-08-1996 19-07-1993 24-06-1993 19-09-1996 09-01-1997 14-10-1996

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/21461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9312108 A		EP 0618913 A	12-10-1994
		ES 2090956 T	16-10-1996
		GR 3021464 T	31-01-1997
		HK 31397 A	21-03-1997
		JP 7121939 B	25-12-1995
		JP 6511009 T	08-12-1994
		KR 142802 B	15-07-1998
		MX 9207393 A	30-06-1994
		NZ 246441 A	27-02-1996
		SG 44510 A	19-12-1997
		US 5463074 A	31-10-1995
		ZA 9209788 A	22-06-1993